

Estimating Regression Models in Which the Dependent Variable Is Based on Estimates

Jeffrey B. Lewis and Drew A. Linzer

*Department of Political Science, University of California,
Los Angeles, 4289 Bunche Hall, Los Angeles, CA 90095
e-mail: jblewis@ucla.edu (corresponding author)
e-mail: dlinzer@ucla.edu*

Researchers often use as dependent variables quantities estimated from auxiliary data sets. Estimated dependent variable (EDV) models arise, for example, in studies where counties or states are the units of analysis and the dependent variable is an estimated mean, proportion, or regression coefficient. Scholars fitting EDV models have generally recognized that variation in the sampling variance of the observations on the dependent variable will induce heteroscedasticity. We show that the most common approach to this problem, weighted least squares, will usually lead to inefficient estimates and underestimated standard errors. In many cases, OLS with White's or Efron heteroscedastic consistent standard errors yields better results. We also suggest two simple alternative FGLS approaches that are more efficient and yield consistent standard error estimates. Finally, we apply the various alternative estimators to a replication of Cohen's (2004) cross-national study of presidential approval.

1 Introduction

It is not uncommon for regressions to be run in which observations on the dependent variable are estimated in an auxiliary analysis. For example, such dependent variables may be calculated as sample means, vectors of regression coefficients, poll question marginals, election vote shares, legislative seat shares, or fractions of bills vetoed; all such variables are “estimated” in the sense that they are aggregated outcomes of individual-level data generating processes. The fitting of such estimated dependent variable (EDV) regressions is the subject of this paper.

In general, EDV regression models are the second stage in a two-stage estimation process. The first stage uses observed data to estimate the values of the dependent variable; the EDV model then regresses these values against one or more independent variables to generate the ultimate coefficients of interest. An obvious context in which EDV regression can arise is multilevel modeling.¹ To foreshadow our application of an EDV model later in this article, consider a simple two-level model in which the individual-level effect of

Authors' note: We thank Chris Achen, Barry Burden, Michael Herron, Gary King, Eduardo Leoni, and Lynn Vavreck for comments on earlier drafts of this article. Any remaining errors are ours alone.

¹Guidance on multilevel models is often derived from such well-cited sources as Steenbergen and Jones (2002) and Bryk and Raudenbush (1992).

evaluations of the economy on leader approval (bottom level) is assumed to be a function of state-level factors plus a random shock (top level). In this manner, a researcher can investigate sources of cross-national causal heterogeneity and address questions of *why* some independent variable may have a different effect on the dependent variable in different countries.

Aside from a loss of efficiency, the fact that the dependent variable is estimated does not necessarily present any difficulties for regression analysis. Indeed, such errors of measurement are often explicitly included in discussions of regression residuals presented in introductory textbooks (Maddala 2001, p. 64). However, if the sampling uncertainty in the dependent variable is not constant across observations, the regression errors will be heteroscedastic and ordinary least squares (OLS) will introduce further inefficiency and may produce inconsistent standard error estimates.

When the dependent variable in a regression is based on estimates, the regression residual can be thought of as having two components. The first component is sampling error (the difference between the true value of the dependent variable and its estimated value). The second component is the random shock that would have obtained even if the dependent variable were directly observed as opposed to estimated. The first component will be heteroscedastic if the sampling variance differs across observations.² But the second component could well be homoscedastic. The usual approaches to EDV regression are either to run OLS ignoring heteroscedasticity resulting from the first component of the residual or to run a weighted least squares (WLS) model assuming that the entire residual, and not just the first component, is heteroscedastic. Under this assumption, however, the WLS approach, like the OLS approach, is also inefficient and may produce inconsistent estimates of parameter uncertainty.

Our main finding for researchers interested in the application of EDV regression models is that only when the share of the regression residual due to sampling error in the dependent variable is *very high* (roughly 80% to 90% or more) does the standard WLS, in which the observations are weighted by the inverse of the standard errors of the dependent variable estimates, produce more efficient estimates than OLS. That is to say, the “cure” provided by the standard WLS is typically going to do more harm than good. Through a series of Monte Carlo experiments, we find that when the sampling uncertainty component is small relative to the total variation in the dependent variable and when this uncertainty does not vary greatly from observation to observation, White’s (1980), and for smaller samples, Efron heteroscedastic consistent standard error estimates yield good results.³ If sampling error comprises a larger share of the variation in the dependent variable and this uncertainty varies greatly across observations, appreciable gains in efficiency can be achieved through the use of either of a pair of alternative feasible generalized least squares (FGLS) estimators that we describe below.

The article is organized as follows. Section 2 reviews applications of models using estimated dependent variables. Section 3 formalizes the problem of regressions involving

²For example, reported average state income may be based on a much larger sample in California than it is in Rhode Island, leading California’s mean income to be more precisely estimated than Rhode Island’s. Some surveys, such as the 1988 Senate Election Study, intentionally draw samples of roughly equal size from each aggregate unit, thus avoiding much of the heteroscedasticity that is generally present when sample means are used as a dependent variable. In these cases, the heteroscedasticity from sampling error would only enter from inter-unit heterogeneity in the intra-unit variance of the variable being sampled.

³Efron standard errors (also known as HC3 standard errors) are based upon the jackknife techniques of Efron (1982) and are typically more accurate (as well as more conservative) than Huber-White standard errors in samples smaller than 250 observations; see Long and Ervin (2000) and MacKinnon and White (1985).

estimated dependent variables. In Section 4 we compare the estimation of EDV models with OLS or WLS to two FGLS estimators that allow the analyst to address the problem of heteroscedasticity in the first-level error component without assuming that the second-level error component is similarly heteroscedastic. The first FGLS approach assumes that the sampling variances of the observations on the dependent variable are known, as would be the case if, for example, the data were regression estimates or aggregated survey responses. The second estimator requires only that the sampling errors be known up to a proportional factor, as would be the case if the data were means based on samples of differing sizes.⁴ Section 5 describes the Monte Carlo experiments. Section 6 presents an empirical example based on the cross-national public opinion study by Cohen (2004). Section 7 concludes.

2 Applications of EDV Models

Multilevel models may be estimated in several ways: (1) The parameters of each of the levels can be estimated simultaneously; (2) Regressions can be run with a large number of interaction terms between within- and across-country variables on the right hand side; or (3) Two-stage approaches of the sort we consider here can be applied (for examples of the first two approaches, see Lubbers et al. 2002; Anderson and Tverdova 2003; Banducci et al. 2003; Peffley and Rohrschneider 2003; and Rohrschneider and Whitefield 2004). Reviewing these alternatives, the “interaction term” model is generally not advised because it assumes, almost certainly incorrectly, that there would be no residual in a regression of individual-level coefficients on the country-level variables. Simultaneous (or single-stage) estimation may take the form of a linear hierarchical model, for example.

Under a two-stage approach, the bottom-level parameters are first estimated within each unit; then, in a second stage, an EDV regression is run in which the estimated quantities from the first stage are used as dependent variables. An example of this multilevel, two-stage modeling strategy can be found in Guerin et al. (2001), who examine determinants of recycling behavior in 15 EU countries using survey data from the Eurobarometer: “The first step was to estimate the person-level model. Then, we set up our model to predict the variation of the coefficients β_{0j} by introducing explanatory variables at the country level” (p. 204). Here, it is regression coefficients as estimates of the relationship at the micro level that become the dependent variable in the top-level cross-national analysis.

The first-stage estimates that become the dependent variable in the second stage need not be regression coefficients—indeed, they may be as simple as aggregate public opinion poll marginals. One such study, which we shall replicate and extend below, is Cohen (2004). He examines country-level approval of the chief executive as a function of each country’s aggregate perceptions of the national economy in 41 countries, with survey sample sizes ranging from 500 to 2189. Because both the estimates *and* the sample sizes differ by country, the standard errors of the country-level dependent variable estimates will also vary by country. Kaltenthaler and Anderson’s (2001) cross-national study of the determinants of aggregate public support for the common European currency is a similar example of using survey marginals as a dependent variable. It is also common to encounter regressions using poll question marginals (such as presidential approval) as dependent

⁴In this case, if the within-unit variance of the variable were constant, the sampling variances would be proportional to $1/n_i$, where n_i is the size of the sample from which the mean was calculated for observation i .

variables in opinion research that is not cross-national (Taylor 1980; MacKuen et al. 1989; MacKuen et al. 1992; Oppenheimer 1996).⁵

While Hanushek (1974) and Jusko and Shively (2005) show that such two-stage approaches can be efficient under certain circumstances and assumptions, in general, there are potential efficiency gains from estimating both the bottom- and top-level parameters in a single stage. In cases in which the amount of information available to estimate the bottom-level effects in each top-level unit is small (particularly relative to the number of top-level units), there are potential gains from “borrowing strength” across units and increasing efficiency through a single-stage estimation that justify their use.

On the other hand, in many political science examples, considerable information is available about the bottom-level effects. In these cases, the gains from estimating both levels in a single step can be modest. To see this, note that no single-step estimator can generate more efficient estimates of the top-level parameters than could be achieved if the bottom-level parameters were known. Taking the simplest example in which the sampling uncertainty in the bottom-level quantities of interest are constant across all top-level units, the standard errors in the second-stage EDV regression are increased by

$$100\sqrt{\frac{S}{1-R^2}}$$

percent over what they would be if the first-level parameters were known, where S is the ratio of estimation variance to all other variance in the second-stage dependent variable and R^2 is the R^2 of the second-stage regression that would obtain if the values of the dependent variable were not based on estimates. In the empirical example presented in Section 6 below, S is approximately 0.005 and R^2 is approximately 0.5. Thus the upper bound on the loss of efficiency that arises from running a second-stage EDV regression is on the order of 10%. Gains from a single-stage estimation would be less than this theoretical upper bound and could be much less. Under these circumstances, the added effort of fitting a more complex single-stage linear hierarchical model would provide little advantage relative to the simple two-stage EDV method.

3 Regressions with Estimated Dependent Variables

Consider the following regression model,

$$y_i = \beta_1 + \sum_{j=2}^k \beta_j x_{ij} + \epsilon_i \quad (1)$$

⁵This very short summary hardly scratches the surface of the breadth and depth of possible applications of EDV regression models. However, we do wish to highlight that with respect to EDVs generated using King’s (1997) EI algorithm, Herron and Shotts (2003) point out that using the so-called precinct-level EI estimates as dependent variables in second-stage regressions will lead to attenuated estimates and more generally calls into question the validity of using precinct-level EI estimates in subsequent analysis. It should be noted the techniques we present below are predicated on the assumption that the data used are free of the features described by Herron and Shotts. In particular, we assume that the sampling or measurement error in the dependent variable ($Y^* - Y$) is independent of the independent variables and error term (\mathbf{X} and ϵ) of the regression. Also, using *district-level* EI estimates (for example, estimates of black turnout at the Congressional district level made by applying EI to precinct-level data in each district) need not involve the same “logical inconsistency” identified by Herron and Shotts.

for observations $i = 1, \dots, N$. Assume that this regression model meets all of the usual Gauss-Markov assumptions. However, y_i is not observable. Rather we observe an unbiased estimate y_i^* where

$$y_i^* = y_i + u_i \tag{2}$$

and $E(u_i) = 0$ and $\text{Var}(u_i) = \omega_i^2$ (for $i = 1, \dots, N$). u_i is the sampling error in y_i^* and ω_i^2 is the variance of that sampling error. Assume that the estimate of the dependent variable for each observation is independent of the others.⁶ In particular, assume $\text{Cov}(u_i, u_h) = 0$ for all $i \neq h$. Now consider the regression formed by substituting Eq. (2) into Eq. (1),

$$y_i^* = \beta_1 + \sum_{j=2}^k \beta_j x_{ij} + u_i + \epsilon_i.$$

Writing $v_i = \epsilon_i + u_i$, we have

$$y_i^* = \beta_1 + \sum_{j=2}^k \beta_j x_{ij} + v_i. \tag{3}$$

Obviously, if $\omega_i \neq \omega_h$ for some i and h , v_i is heteroskedastic. In particular, letting $\mathbf{v} = (v_1, v_2, \dots, v_N)'$,

$$E(\mathbf{v}\mathbf{v}') = \mathbf{\Omega} = \begin{bmatrix} \sigma^2 + \omega_1^2 & 0 & \dots & 0 \\ 0 & \sigma^2 + \omega_2^2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \dots & \dots & \sigma^2 + \omega_N^2 \end{bmatrix} \tag{4}$$

where $\text{Var}(\epsilon) = \sigma^2$. If σ^2 and ω_i^2 were both known, a WLS approach would yield best linear estimators. That is, OLS estimation of

$$w_i y_i = \beta_1 w_i + \sum_{j=2}^k \beta_j x_{ij} w_i + w_i v_i$$

where

$$w_i = \frac{1}{\sqrt{\sigma^2 + \omega_i^2}} \tag{5}$$

yields efficient estimates of the regression parameters.

In real-world applications, we generally have only estimates of the ω_i^2 ($i = 1, \dots, N$) and no knowledge of σ^2 . In the next section, we consider four feasible approaches to this problem in cases in which knowledge of σ^2 and/or the ω_i^2 's is missing or incomplete.

4 Approaches to Fitting Estimated Dependent Variable Regressions

We will consider four approaches to estimating EDV regressions. The first two are the most commonly applied: OLS (with or without White's or Efron heteroscedastic consistent standard errors) and WLS using weights based on the inverse of the standard errors of the

⁶The first FGLS approach described below is trivially extended to the case in which the estimates are not independent.

dependent variable estimates. The third is an FGLS procedure first suggested in Hanushek (1974) appropriate for cases in which estimates or the true sampling variances of the y_i^* are available (i.e., the ω^2 's are known). The fourth is also an FGLS procedure appropriate for cases in which the sampling variances of the y_i^* 's are known only up to a constant of proportionality. Both of these estimators are very easy to implement using standard statistical packages.⁷

4.1 Estimation by Ordinary Least Squares

Given the assumptions made previously, it is well known that Eq. (3) can be consistently estimated by OLS. However, in order for OLS to be efficient, ω_i must be constant across all observations. That is, only if the sampling variances of the y^* are constant (or zero) will v_i be homoscedastic and OLS efficient.

In general, OLS will not be efficient and the usual standard error estimator is, in general, inconsistent. Thus OLS estimates will be less precise than is optimal and, under some conditions, will produce badly inconsistent standard error estimates. Not surprisingly, if the degree of variation in ω_i^2 is small or if σ^2 is large relative to the ω^2 's, then OLS will perform quite well. That is, if the appropriate set of weights (w_i for $i = 1, \dots, N$) are nearly constant across observations, OLS will be nearly efficient.

The inconsistent OLS standard errors can be corrected using the White (1980) or Efron (1982) heteroscedastic consistent standard error estimator. As will be shown below, the Efron robust standard error estimator will correctly measure the uncertainty in OLS even in fairly small samples. However, the OLS estimator may be quite inefficient in some cases. OLS is inefficient because the partial information about the nature of the heteroscedasticity (that is, knowledge of ω_i^2 's) is not used.

4.2 Estimation by Weighted Least Squares

The most common WLS approach to EDV models is simply to set $w_i = 1/\omega_i$ ($i = 1, \dots, N$). Such weighting is recommended in Saxonhouse (1976) and King (1997, p. 290), among other places. As shown in Eq. (5), this weighting scheme implicitly sets $\sigma^2 = 0$. In other words, when weighting by $1/\omega$ in an EDV model, researchers are attributing *all* of the error in the total residual v_i to the heteroscedastic sampling error (u_i) and none to the homoscedastic noise (ϵ_i). This is equivalent to assuming that the R^2 in their regression would be 1 ($\sigma^2 = 0$) if they directly observed the true y 's as opposed to the estimated y^* 's.

Of course, the assumption that the regression R^2 would be 1 if the true dependent variable was directly observable will not typically be met. However, as will be shown in the Monte Carlo simulation in the next section, if the variance of the homoscedastic noise (σ^2) is very small in comparison to the variances of the sampling errors (the ω_i^2 's), the assumption that $\sigma^2 = 0$ may not be too costly. However, as σ^2 grows relative to the ω_i^2 's, the WLS estimator becomes increasingly—and quite badly—inefficient and generates highly misleading standard error estimates.⁸

Given only these two options, researchers face a trade-off. Use OLS with robust standard error estimates and get possibly inefficient parameter estimates but good estimates of the parameter uncertainty. Or use WLS and get possibly much more efficient parameter estimates, but risk getting very misleading standard errors. Some scholars have,

⁷R functions implementing both of these procedures are available from the authors.

⁸These conclusions are not specific only to the EDV case, but generalize to other cases in which WLS is applied using incorrect weights (see Greene 2003).

in fact, presented both estimates (Burden and Kimball 1998). This is a good practice. However, when the methods yield estimates that differ in substantively significant ways, simply reporting both sets of results is not entirely satisfying.

In the next two subsections, we will present two alternative estimation techniques that are asymptotically efficient and whose standard errors are consistent regardless of the relative size of σ^2 and the ω^2 's. Both of these estimators are FGLS estimators that use OLS to generate consistent estimates of the v 's. These estimated v 's are used to find consistent estimates of $\omega_i^2 + \sigma^2$ (for $i = 1, \dots, N$) from which weights are created and a second-stage WLS model fit.

4.3 Estimating $\omega^2 + \sigma^2$ when ω^2 is known

In this section, we present an FGLS estimator that is appropriate for cases in which the standard errors of the estimated dependent variable are known, or at least where reliable estimates are available. This method was first suggested by Hanushek (1974) for the case in which the dependent variable is estimated regression coefficients. Since the ω 's are assumed to be known, only an estimate of σ^2 (the variance of the component of the regression residual that is not due to sampling of the dependent variable) is required in order to construct a proper set of weights to use in a second-stage WLS regression. To arrive at an estimate of σ^2 , first run OLS and calculate residuals \hat{v}_i for $i = 1, \dots, N$. Following Hanushek (1974), the expectation of the sum of squared residuals from this OLS regression can be written as

$$E\left(\sum_i \hat{v}_i^2\right) = E(\mathbf{v}'\mathbf{v}) - \text{tr}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{\Omega}\mathbf{X}),$$

where $\mathbf{\Omega}$ is the variance covariance matrix of the vector of regression residuals \mathbf{v} given in Eq. (4) and $\text{tr}(\cdot)$ is the trace operator.⁹ $E(\mathbf{v}'\mathbf{v}) = E(\sum_i v_i^2)$ and

$$\begin{aligned} E\left(\sum_i v_i^2\right) &= E\left[\sum_i (\epsilon_i^2 + u_i^2 + 2\epsilon_i u_i)\right] \\ &= E\left[\sum_i \epsilon_i^2\right] + E\left[\sum_i u_i^2\right] + 0 \\ &= N\sigma^2 + \sum_i \omega_i^2. \end{aligned}$$

Writing $\mathbf{\Omega}$ as $\sigma^2\mathbf{I} + \mathbf{G}$ where \mathbf{I} is an $n \times n$ identity matrix and \mathbf{G} is an $n \times n$ diagonal matrix with ω_i^2 as the i th diagonal element, we have

$$\begin{aligned} E\left(\sum_i \hat{v}_i^2\right) &= N\sigma^2 + \sum_i \omega_i^2 - \text{tr}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'(\sigma^2\mathbf{I} + \mathbf{G})\mathbf{X}) \\ &= N\sigma^2 + \sum_i \omega_i^2 - \sigma^2\text{tr}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}) - \text{tr}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{G}\mathbf{X}) \\ &= (N - k)\sigma^2 + \sum_i \omega_i^2 - \text{tr}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{G}\mathbf{X}), \end{aligned}$$

⁹The trace of a square matrix is the sum of its diagonal elements.

where k is the number of columns in \mathbf{X} . Rearranging, we have

$$\sigma^2 = \frac{E(\sum_i \hat{v}_i^2) - \sum_i \omega_i^2 + \text{tr}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{G}\mathbf{X})}{N - k},$$

implying that

$$\hat{\sigma}^2 = \frac{\sum_i \hat{v}_i^2 - \sum_i \omega_i^2 + \text{tr}((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{G}\mathbf{X})}{N - k}$$

is an unbiased estimator for σ^2 . In small samples, σ^2 may be estimated to be less than 0. In these cases, σ^2 can be set equal to 0 (yielding the standard WLS approach to EDV regression).

Given this estimator for σ^2 , a set of weights w_i (for $i = 1, \dots, N$) can be constructed such that

$$w_i = \frac{1}{\sqrt{\omega_i^2 + \hat{\sigma}^2}}.$$

The main regression is then refit using these weights. Estimates from this regression will be asymptotically efficient.

4.4 Estimating $\omega^2 + \sigma^2$ when ω^2 is known to a constant multiplicative factor

Sometimes researchers encounter situations in which they may know only the ω^2 's up to some unknown constant of proportionality. A common example here would be a dependent variable based on sample means. That is, $y_i^* = \sum_{m=1}^{n_i} Y_{im}/n_i$. Often, in such cases, we know the size of the samples upon which the estimates are based, but we do not know the exact sampling variances of the estimated means. If we assume that the variance of Y_{im} is constant for all i and j , then we have

$$\omega_i^2 = \frac{\kappa}{n_i},$$

where κ is the variance of Y and n_i is the size of the sample upon which the i th sample mean is based.¹⁰

¹⁰The usual WLS approach described above is often advocated for this case (Hanushek and Jackson 1977). The justification is as follows. Suppose that all the variables are measured as sample means. Then assume there is an underlying individual-level regression model,

$$Y_{im} = \beta_1 + \sum_{j=2}^k \beta_j x_{imj} + \epsilon_{im}.$$

Assuming that ϵ_{im} is i.i.d. and taking means within each sampling unit i , we find,

$$y_i^* = \beta_1 + \sum_{j=2}^k \beta_j \bar{x}_{ij} + \bar{\epsilon}_i,$$

where the variance of $\bar{\epsilon}_i$ will be proportional to n_i (the number of individuals sampled in each unit i). In this case, weighting the entire regression residual by $\sqrt{n_i}$ would be appropriate. However, if some x 's that are uncorrelated to ϵ at both the individual and aggregate level are omitted, we will have an additional error component that is not heteroscedastic in proportion to $1/n_i$.

Again, we first run OLS and use the estimated residuals to form a set of weights, w_i ($i = 1, \dots, N$) that can then be used in a second-stage WLS regression. We begin by noting that the expectation of the true squared error terms is

$$E(v_i^2) = \sigma^2 + \frac{\kappa}{n_i}.$$

If $\mathbf{v} = (v_1, \dots, v_N)'$ were observed directly, σ^2 and κ could be estimated efficiently by the OLS regression of v^2 on $(1/n)$. Since the \hat{v} 's are consistent estimators for the v 's, we have

$$\text{plim}_{N \rightarrow \infty} E(\hat{v}_i^2) = \sigma^2 + \kappa \left(\frac{1}{n_i} \right).$$

One can then consistently estimate σ^2 and κ by regressing \hat{v} on $(1/n)$. As above, this method can yield estimates of $\hat{\sigma}^2 < 0$. In these cases, the regression of \hat{v}^2 on $(1/n)$ can be rerun constraining the constant ($\hat{\sigma}^2$) to be 0. Predicted values from the regression of \hat{v}^2 on $(1/n)$ are estimates of $\sigma^2 + \omega_i^2$. Using these estimates, we next construct a set of weights where

$$w_i = \frac{1}{\sqrt{\hat{\sigma}^2 + \hat{\kappa}(1/n_i)}}.$$

The general procedure is as follows:

1. Regress \mathbf{y}^* on \mathbf{X} by OLS and calculate the squared residuals \hat{v}_i^2 for $i = 1, \dots, N$.
2. Regress \hat{v}^2 on $\tilde{\omega}_i^2$ where $\tilde{\omega}_i^2$ is proportional to the variance of u (i.e. $\tilde{\omega}_i^2 = \omega_i^2/\kappa$). If the constant in this regression is negative ($\hat{\sigma}^2 < 0$), the regression is rerun constraining the constant to be 0. Calculate predicted values from this regression. These predicted values are consistent estimates of $\omega_i^2 + \sigma^2$.
3. Fit the WLS regression of \mathbf{y}^* on \mathbf{X} using weights $w_i = \frac{1}{\sqrt{\hat{\sigma}^2 + \hat{\kappa}\omega_i^2}}$.

Because this method involves estimating not only σ^2 , but also κ , it will yield less-efficient estimates than those that can be achieved if the ω^2 's are known exactly and only σ^2 must be estimated.

5 Monte Carlo Analysis

In this section, we describe a series of Monte Carlo experiments that we conducted in order to ascertain the small sample properties of each of the estimators for the EDV model described above.

The regression model used for all the experiments is

$$y_i = 1 + 1x_i + v_i,$$

where $v_i = \epsilon_i + u_i$. Both ϵ and u are drawn from independent (conditional) normal distributions, $\epsilon_i \sim N(0, \omega_i^2)$ and $u_i \sim N(0, 1 - C)$ for $i = 1, \dots, N$. The ω^2 's are drawn from a gamma distribution. In particular, $\omega_i^2 \sim \text{Gamma}(C/\theta, 1/\theta)$.¹¹

¹¹The parameterization of the gamma distribution used here follows DeGroot and Schervish (2002). We define the density of gamma distribution as $f(z | \alpha, \beta) = [\Gamma(\alpha)]^{-1} \beta^\alpha z^{\alpha-1} e^{-\beta z}$ for $\alpha > 0$, $\beta > 0$, and $z > 0$. Given this parameterization, $E(Z) = \alpha/\beta$ and $\text{Var}(Z) = \alpha/\beta^2$. In the simulations, the density of ω_i^2 is $f(\omega_i^2 | C/\theta, 1/\theta)$.




Fig. 1 Shows the distributions of ω_i^2 s (sampling variances of the observations on the dependent variable) in the Monte Carlo experiments. Notice that as θ gets smaller, the frequency of extreme outliers decreases.

Letting u be distributed as a mixture of normals where the normal distribution’s variance parameter is distributed gamma is convenient because, as shown by Patel and Read (1996, p. 31), the variance of w_i^2 will be

$$\text{Var}(\omega_i^2) = C\theta.$$

Therefore, the total variance of u_i (without conditioning on ω_i^2) is

$$\text{Var}(u_i) = \frac{C/\theta}{1/\theta} = C.$$

Thus, across the entire sample, $\text{Var}(v) = 1$ and C represents the share of the total regression error that is due to sampling error in the dependent variable.¹² The variance of X is also set to 1, so the R^2 in the regressions is approximately 0.5 in all cases.¹³ The parameter θ describes the degree of dispersion in the ω^2 s across observations. As shown in Fig. 1, the larger θ is, the more dispersed are the ω^2 s. That is, the larger θ is, the more variation there is in the accuracy with which the dependent variable is estimated across observations. As θ approaches 0, the distribution of ω_i collapses to a point. As noted above, if ω_i is constant across observations, the error term in the regression will not be heteroscedastic and OLS regression will be efficient.

The parameters, the features of the data they relate to, and values they are set to in the experiments are shown in Table 1. Before describing the results, we will first review our expectations about how each of the parameters will affect the four estimators for the EDV model developed in Section 3.

When C is close to zero, sampling error in the dependent variable makes only a small contribution to the overall residual. In this case, the overall residual will be nearly homoscedastic and OLS should perform quite well. On the other hand, a WLS regression

¹²Note that $v_i = u_i + \epsilon_i$. Because ϵ_i is assumed to be independent of u_i , $\text{Var}(v_i) = \text{Var}(u_i) + \text{Var}(\epsilon_i) = C + (1 - C) = 1$.

¹³The “explained” variance will be $\beta^2\text{Var}(X) = 1$ and the “unexplained” variance $\text{Var}(v) = 1$, thus the R^2 will be approximately $1/(1 + 1) = 1/2$.

Table 1 Parameters for the Monte Carlo experiments

<i>Parameter</i>	<i>Description</i>	<i>Values</i>
C	Share of variation in the regression error due to estimation error in the dependent variable	(0.1, 0.2, . . . , 1.0)
θ	Degree of variation in dependent variable sampling variances across observations	(0.2, 0.5, 0.8)
ρ	Correlation between X and sampling variances of the dependent variable	(0.0, 0.5, 1.0)
N	Number of observations	30
β	Regression slope	1

Note. Table describes parameters that were manipulated in the Monte Carlo experiments and the values to which they were set. For each combination of parameter values, 80,000 simulations were performed.

that weights by $1/\omega$ will “overcorrect” the heteroscedasticity that is generated by estimation errors in the dependent variable. Thus, *ceteris paribus*, low values of C should lead to poor WLS estimates.

The larger θ is, the more heteroscedasticity is introduced by the sampling errors in the dependent variable. When θ is small, we expect that OLS will produce good estimates because the regression error will exhibit little heteroscedasticity even if C is large. However, if θ is large, OLS is expected to be inefficient, particularly as C increases.

The value of ρ is mainly expected to affect the accuracy of the standard errors. It is well known that OLS will produce inconsistent standard error estimates only if the heteroscedasticity of the regression error depends on the independent variables in some way. Thus we expect that OLS will be relatively efficient if ρ is small. However, as ρ grows, the efficiency of the OLS estimates should decline and the standard error estimates worsen.

OLS combined with White’s (1980) standard errors should produce reasonable estimates of model uncertainty for all parameter settings though the estimates may be inefficient. The FGLS estimators should be efficient and produce good standard error estimates for all parameter values.¹⁴

In all of the experiments described previously, we use data sets with 30 observations. Similar results are obtained in experiments using data sets with greater numbers of observations. For each combination of the parameter values, 80,000 simulations were performed.

Table 2 shows the results for one particular set of parameters. In this experiment, $C = 0.7$, meaning that 70% of the variance in the residual is due to sampling error in the dependent variable. The variation in the sampling error of the dependent variable across observations is high ($\theta = 0.8$). In particular, across the simulations the largest sampling variance was about 200 times as large as the smallest on average. The correlation between the sampling variances and the independent variable is set to 0.5.

All of the estimators yield estimates that are very accurate on average. Across the simulated data sets, the average estimated regression slope was nearly identical (to four decimal places) to the true value of β . This is not surprising, given that all the estimators tested are known to be consistent, but it is comforting that they show little to no bias in samples as small as 30. This is generally true for all of the combinations of parameter values given in Table 1. Indeed, across all of the experiments that we conducted the mean $\hat{\beta}$

¹⁴Exceptions to this claim are $C = 0$ and $C = 1$, where OLS and WLS, respectively, would be efficient.

Table 2 Monte Carlo results for various EDV estimators

<i>Method</i>	<i>Regression slope ($\hat{\beta}$)</i>			
	<i>Mean</i>	<i>Observed root mean square error (RMSE)</i>	<i>Mean estimated standard error (SE)</i>	<i>RMSE as a percentage of mean est. SE</i>
OLS				
Usual SEs	1.00	0.211	0.188	112
White's SEs	1.00	0.211	0.184	115
Efron SEs	1.00	0.211	0.214	98
WLS	1.00	0.228	0.158	144
FGLS				
Known variance	1.00	0.189	0.174	109
Proportional variance	1.00	0.193	0.171	113

Note. Shows the results of one Monte Carlo experiment described in the text. The numbers reported in the table are based on 200,000 simulations, with $N = 30$ observations each. The parameter values are: $\beta = 1$, $\rho = 0.5$, $\theta = 0.8$, and $C = 0.7$. Even when 70% of the regression residual is due to estimation error in the dependent variable, the usual WLS approach is much less efficient than OLS or the alternative FGLS approaches. The observed standard error of $\hat{\beta}$ is larger using WLS than it is using OLS. Moreover, WLS and, to a lesser extent, OLS systematically underestimate the uncertainty in $\hat{\beta}$.

did not differ from the true β in a statistically significant way. One convenient consequence of all the estimators showing little to no bias is that their observed root mean square errors (RMSEs) are basically equivalent to their observed sampling standard deviations. Thus the observed RMSEs can be compared to the mean estimated standard errors to assess the degree to which each model accurately represents the uncertainty of its estimates.

While the parameter values used to generate the results in Table 2 would seem to approximate the conditions under which the WLS approach would be effective, the experimental results suggest otherwise. Even with very highly dispersed ω^2 's and most of the residual resulting from sampling error in the dependent variable, OLS produced more efficient estimates with more accurate standard errors than did the standard WLS. The FGLS estimators are both about 10% more efficient than OLS; further experimentation (not reported here) indicates that the efficiency gain from the FGLS models is even greater in larger sample sizes. Additionally, both of these FGLS estimators produced accurate standard error estimates in this experiment.

WLS produced highly misleading standard error estimates. While the standard deviation of the WLS slope estimator across the repetitions was 0.228 (see the "RMSE" column in the table), the average standard error estimate was 0.158. Thus, on average, the observed standard deviation of the estimated β across the simulations was 144% greater than the estimated standard errors. As we expected, OLS also produces inconsistent standard error estimates. However, the observed OLS root mean square error was only 112% greater than the mean estimated OLS standard error. White's (1980) heteroscedastic consistent standard error estimator actually does not improve the estimation of the standard error of β a great deal in this example. However, accurate (albeit somewhat conservative) estimates of OLS parameter uncertainty were obtained using Efron small-sample standard error estimates. As noted earlier, as the size of the sample increases, White's standard errors will increasingly outperform the usual OLS standard errors as well as the Efron standard errors. Finally, following our expectation, both of the FGLS estimators produced good estimates of parameter uncertainty.

Overall, in this experiment, WLS was clearly inferior to OLS and the FGLS estimators. Even with 70% of the total regression error resulting from sampling error in the dependent variable and vast dispersion in the sampling variances across observations, OLS is preferable to the standard WLS approach. OLS with Efron standard errors produced quite satisfactory results. However, the FGLS estimators, which use the information about the sampling errors in the dependent variable, did produce percent efficiency gains over OLS. Whether these same conclusions hold more generally is the question to which we now turn.

Figure 2 graphs the observed standard deviations of the estimates (RMSEs) as a function of the percent of the total regression error that is due to sampling the dependent variable for various values of θ and ρ . The results are consistent with expectations. When θ is low—when variance of the sampling errors in the dependent variable is fairly constant across observations—all of the methods perform very similarly. Indeed, the lines representing each of the methods in the top three panels of the graph are difficult to distinguish. As the variation in the dependent variable sampling errors across observations (θ) increases, differences in the behavior of the estimators becomes apparent. In particular, when θ is large and little of the total regression residual stems from the mismeasurement in the dependent variable (C is small), WLS produces estimates with approximately twice the RMSE as OLS. Even as this fraction is increased, OLS continues to outperform WLS. Indeed, until about 80% of the total error variance is the result of sampling error in the dependent variable, OLS produces more efficient estimates than does the standard WLS approach.¹⁵

As anticipated, the FGLS estimators produce efficient estimates relative to OLS and WLS, though in many cases the gains are quite modest. The “known variance” FGLS estimator—the estimator that requires that the sampling variances of the dependent variable observations be known—is generally more efficient than the proportional variance estimator. This result follows from the fact that the proportional variance estimator requires the estimation of one more parameter than does the known variance estimator.

Figure 3 graphs the observed standard errors across the 80,000 experiments as a fraction of the average estimated standard errors of the various estimators against the percent of the total regression error that is due to sampling error in the dependent variable. Overall, these graphs are consistent with the expectations laid out earlier. OLS tends to produce biased standard error estimates for high values of C , and WLS produces biased standard error estimates except when C is very high (over 0.9). White’s standard error estimates are generally effective, but in small samples, if robust standard errors are to be employed, we recommend the more conservative Efron standard errors. With small $n = 30$ in these simulations, the Efron estimates are more consistently accurate under each of the nine scenarios considered. The two FGLS estimators produce quite accurate estimates of parameter uncertainty; the known variance FGLS estimator works particularly well in all cases. The proportional variance FGLS estimator is less successful when C is low, below 0.4, but generally performs no worse than WLS, especially with high values of θ .

Moving from the left-most to right-most panels of Fig. 3, one sees that as the correlation between the independent variable and the sampling variances of the dependent variable increases the bias of the OLS standard errors also increases.¹⁶ However, the Efron robust standard errors are not similarly affected. It is interesting to note that even in the low sampling error dispersion case ($\theta = 0.2$), the observed OLS and WLS standard errors are in some cases as much as 110% to 120% greater than the estimated standard errors. While

¹⁵When we ran the same simulation with a larger $n = 500$, OLS was more efficient than WLS until about 90% of the total error variance was the result of sampling error in the dependent variable.

¹⁶This result is similar to those typically found in the heteroscedasticity literature (Greene 2003, p. 505).




Fig. 2 Shows the root mean square error of the estimated regression slope coefficient. Notice that WLS can be particularly inefficient when very little of the regression residual is due to estimation error in the dependent variable. Simulations are for 80,000 samples of $n = 30$ observations.

all the methods are quite similar in terms of their efficiency when $\theta = 0.2$ (see Fig. 2), the FGLS estimators or OLS with Efron standard errors are still preferable because they result in better estimates of parameter uncertainty.

6 Empirical Application: Economic Perceptions and Leader Approval

Do individuals evaluate their political leaders based on prospective or retrospective assessments of the economy? Does this relationship depend on whether or not individuals reside in a country that is democratic and economically advanced? These are the questions addressed by Cohen (2004) in a recent cross-national public opinion study, in response to a debate initiated by MacKuen et al. (1992) and continued by Clarke and Stewart (1994) and Norpoth (1996) over the determinants of approval ratings of the U.S. president. Cohen




Fig. 3 Shows the observed standard errors of the regression slope as a percentage of its average estimated standard error. Notice that under some conditions OLS and WLS standard errors can be very misleading. Simulations are for 80,000 samples of $n = 30$ observations.

argues that “in the advanced industrialized nations of the West, leader approval will be a function of prospective economic perceptions. In contrast, in the developing world and among newer democracies, leader approval will be a function of retrospections” (2004, p. 31). To conduct his analysis, Cohen uses survey data from the 2002 Pew Global Attitudes Project, which was fielded between July and October 2002 in 44 countries worldwide, including both developed and developing nations.¹⁷

¹⁷The data set is free to download from the Pew Research Center for the People and the Press data archive at <http://people-press.org/dataarchive/>. See the June 3, 2003, release of the report titled “Views of a Changing World.” The Pew Global Attitudes Project bears no responsibility for the analyses or interpretations of the data presented here.

What makes this study particularly useful for our purposes is that at the time of his analysis, Cohen had access only to the country-level topline that were published in Pew's initial survey report. Therefore, the dependent variable was, by necessity, the *country-level* proportion of respondents approving of their country's president or prime minister. This makes it impossible to draw inferences about the possible relationship between *individuals'* economic perceptions and their approval of their country's chief executive. Since that time, however, Pew has released the full data set of individual-level survey responses. We employ these data to recast Cohen's country-level analysis as a multilevel model in which parameter estimates from within-country, individual-level logit regressions are modeled as a function of economic development at the country level.

Cohen proposes a model that explains country-level presidential approval—"the percentage of respondents in each country who give the leader 'a very good' or 'somewhat good' evaluation" (p. 33)¹⁸—as a function of three variables: economic "prospections," economic "retrospections," and a dummy variable for whether or not a country is an "old" (economically advanced and stable) democracy.¹⁹ Cohen acknowledges the possibility of nonconstant variance in his dependent variable as a result of the survey sample sizes ranging from six countries with 500 respondents to India with 2189 respondents, so he presents his linear regression results with both regular OLS standard errors and robust Huber-White corrected standard errors.

Though Cohen does not present a weighted least squares (WLS) model in his results, one could be easily estimated. Because the survey responses to the question of leader approval arise from a random sample, we could estimate the standard error of that estimate as

$$\widehat{SE}(P_i) = \frac{\sqrt{P_i(100 - P_i)}}{n_i},$$

where P_i is the percent support for the president or prime minister in a given country and n_i is the number of survey responses in country i . The inverse of these estimates would then become the weights in a WLS regression. Looking more carefully at the standard errors of the presidential approval ratings across the 41 surveyed countries in this example highlights the concern associated with the application of a WLS model. Between the countries with large and small sample sizes, the standard error estimates range only from a low of 0.77 (Uzbekistan) to a high of 2.23 (Canada). This disparity is not terribly large—nor is it ever likely to be for cross-national public opinion studies in which the difference in the number of interviews conducted in different countries will rarely exceed one order of magnitude. In the notation of the Monte Carlo results presented earlier, this is a case in which the variation in the sampling variances across observations, θ , is small to moderate.

A good gauge of whether WLS is appropriate in this case is the R^2 that the regression would achieve if the only source of error was the uncertainty due to estimation of the dependent variable. This R^2 can be approximated as one minus the average sampling variance of Y^* (the residual variance) over the observed variance of Y^* across countries (the total variance).²⁰ In this case, the average estimated sampling variance is about 2.8, while the

¹⁸The approval rating question—question 35b in the survey—was asked in 41 of the 44 study countries: all except China, Egypt, and Vietnam.

¹⁹The economic retrospection and prospection questions are numbers 12 and 13, respectively. The model includes two interaction terms to capture the possibility that the effect of economic evaluations differs between countries that are "old" and those that are not.

²⁰The approximate $R^2 = 1 - \frac{0\bar{r}^2}{S_{y,*}^2}$ where $S_{y,*}^2$ is the sample variance of the observed dependent variable across the observations on the dependent variable.

cross-country variance of leader support is 513.7, so the R^2 that one would expect is greater than 0.99.²¹ Cohen reports an R^2 from his OLS regression Model 3 (p. 35) to be a much more modest 0.51. Thus, of the roughly 49% of the variance in leader approval that is not accounted for by his model, only about 1% can be attributed to sampling errors in the measurement of the true underlying leader approval. In the notation of the Monte Carlo experiments, C , the fraction of the error due to estimation of the dependent variable, is quite small. As such, WLS estimates are unlikely to be as reliable as the estimates from the OLS model.

With the individual-level data now available from Pew, we reanalyze Cohen's claim that "citizens in advanced democracies employ prospective evaluations, while citizens of less advanced democracies employ retrospective assessments in evaluating the national executive" (pp. 37–38) using a multilevel model that compares the OLS/robust standard error, WLS, and FGLS approaches to EDV regressions.

We first run a series of 41 logistic regression models on the individual-level data within the countries in the Pew study. The dependent variable in these models is whether or not individuals have a "very good" or "somewhat good" opinion of their country's president or prime minister. The independent variables are the individual's responses to the prospective and retrospective economic evaluation questions; both are included in each multivariate model as simultaneous controls. The prospective question is coded across five choices, varying from whether the respondent expects economic conditions over the next 12 months to "improve a lot" (1) all the way to "worsen a lot" (5). The retrospective question is coded across four choices, varying from whether the respondent thinks the current economic situation is "very good" (1) to "very bad" (4). Theory predicts negative coefficients on both variables, indicating that as individuals take an increasingly worse view of their country's economy, they are less likely to support their country's leader. This is indeed what we find: all 41 of the "retrospective" coefficients are negative, as are 39 of the "prospective" coefficients; neither of the remaining two (in Brazil and Jordan) are statistically significant.

Cohen's theory predicts cross-national heterogeneity in these logit coefficient estimates, such that the effect of prospection should be greater, and the effect of retrospection lesser, in countries that are advanced democracies versus those that are not. The coding scheme Cohen devises for the dummy variable for "old" advanced democracies is primarily a function of that country's wealth; the seven countries coded as "old" are the seven wealthiest in the sample.²² We therefore test Cohen's prediction by regressing the within-country coefficient estimates on prospection and retrospection on real per capita GDP in 2000.²³ The results from these second-stage regressions are given in Table 3, for OLS with Efron standard errors, WLS, and FGLS (known variance) models.

Contrary to Cohen's hypothesis, there is *no effect* of wealth on the power of prospective economic considerations to predict leader approval. This finding may be confirmed visually in the left-hand panel of Fig. 4. Individuals are more likely to hold an unfavorable view of their leader as their opinion of the future economy worsens, regardless of the level of wealth of the country in which they reside.

For our purposes, the retrospective models provide a more interesting example. Note first that the sign on the coefficient of log of GDP per capita is opposite what Cohen predicts. If citizens of poorer countries were more heavily economically retrospective when forming opinions of their leader, then the coefficient on wealth would be *positive*,

²¹That is, $0.995 = 1 - (2.8/513.7)$.

²²Cohen codes the "old democracy" variable 1 for Canada, France, Germany, Great Britain, Italy, Japan, and the United States, and 0 otherwise; see also Cohen's footnote 5.

²³These GDP data come from the World Bank 2002 World Development Indicators data set.

Table 3 Testing for cross-national parameter heterogeneity

	<i>Prospective models</i>			<i>Retrospective models</i>		
	<i>OLS</i>		<i>FGLS</i>	<i>OLS</i>		<i>FGLS</i>
	<i>Efron SE</i>	<i>WLS</i>		<i>Efron SE</i>	<i>WLS</i>	
Constant	-0.261 (0.260)	-0.187 (0.254)	-0.221 (0.264)	-0.223 (0.274)	0.057 (0.261)	-0.141 (0.279)
log(per capita GDP)	-0.041 (0.071)	-0.043 (0.068)	-0.046 (0.071)	-0.078 (0.074)	-0.146 (0.070)	-0.098 (0.075)
R^2	0.008			0.026		
$\hat{\sigma}$	0.221		0.189	0.234		0.176
Mean ω			0.109			0.145
N	41	41	41	41	41	41

Note. The dependent variable is the logit coefficient from individual-level regression models in 41 countries. The within-country logit models simultaneously regress leader approval against both prospective and retrospective economic evaluations, thus generating the two sets of 41 coefficients used as the dependent variables here. Data are from countries surveyed in the 2002 Pew Global Attitudes Project and from the World Bank 2002 WDI data set. Standard errors are in parentheses.

not negative. That is, the predicted coefficient on the within-country retrospective variable would be more negative for poorer countries and increase to zero as countries increased in wealth. This is not what we find (see the right-hand panel of Fig. 4). So, from a substantive perspective, we conclude on the basis of this evidence that Cohen’s hypothesis is incorrect.

That said, while the coefficient on the log of GDP per capita is not significantly nonzero in the OLS and FGLS models, the coefficient *is* significant in the WLS model. Why does this difference arise, and which of these estimators is more likely to be producing accurate estimates? Compared to Cohen’s country-level model discussed above, the average estimated sampling error (0.023) is now much closer to the inter-country variance in the retrospection coefficient estimate (0.055). Calculating as before, one would expect an R^2 value of 0.58, but the R^2 from the OLS regression in Table 3 is a mere 0.026. Of the 97% of the variance in the retrospective logit coefficients not accounted for by the model, about




Fig. 4 Scatterplots of logit coefficients on economic prospecting (left) and retrospection (right) from 41 country-level regressions of leader approval on those two variables, versus log of GDP per capita in 2000. Vertical bars are 95% confidence intervals for each coefficient estimate. Best fit lines are given for OLS (solid), WLS (dotted), and FGLS (dashed).

58% of that amount can be attributed to sampling errors in the estimation of the logit coefficients. The remaining nearly 40% of the variance is being erroneously attributed to measurement error by the WLS estimator. This is still a large enough amount that we conclude that the WLS estimator has returned results that are misleading and would lead to a researcher drawing false inferences.

7 Conclusion

Through a series of Monte Carlo experiments, we have demonstrated the following relationships between estimation of EDV regression models using OLS, WLS, and a pair of alternative FGLS estimators:

1. The larger the share of the regression residual that is due to sampling error in the dependent variable, the less efficient OLS is and the more efficient WLS is.
2. WLS leads to greater overconfidence (downwardly biased standard errors) the smaller the fraction of the error variance attributable to sampling error in the dependent variable.
3. OLS standard errors are increasingly biased as the correlation between the independent variables and the variance of the sampling error in the dependent variable increases and as the fraction of the total regression error variance due to sampling error in the dependent variable increases.
4. White's (1980) and Efron (for small samples) heteroscedastic consistent standard error estimators are generally reliable (no over- or under-confidence), though OLS may be quite inefficient.
5. The FGLS estimators presented above produce efficient estimates (relative to OLS) if a sufficiently high fraction of the total regression error variance is due to sampling error. The standard errors of these estimators generally produce less overconfidence than is found using OLS and WLS.

The results presented in this article suggest that information about the variance of the sampling errors in estimated dependent variables should be used with caution. The usual approach of weighting the dependent variable by the inverse of the standard errors of the dependent variable estimates will in most cases lead to inefficient parameter estimates and overconfidence in these estimates. This overconfidence can be very large. In some cases, the true uncertainty in the parameter estimates was nearly double the WLS estimated parameter uncertainty. Discarding information about the sampling errors in the observations on the dependent variable and fitting OLS with White's or Efron robust standard errors is generally superior to the WLS approach. Indeed, OLS with robust standard errors is probably the best approach, except when information about the sampling errors in the dependent variable is not only available, but highly reliable.

However, when reliable information about the sampling variances of the estimated dependent variable is available, the two FGLS approaches presented above will yield generally superior results to OLS. In some of the cases considered, the gain in efficiency was substantial. These FGLS estimators are easy to implement and allow the analyst not only to achieve efficient parameter estimates, but also to estimate the fraction of the total regression error that is due to sampling errors in the measurement of the dependent variable.

References

- Anderson, Christopher J., and Yuliya V. Tverdova. 2003. "Corruption, Political Allegiances, and Attitudes toward Government in Contemporary Democracies." *American Journal of Political Science* 47:91–109.

- Banducci, Susan A., Jeffrey A. Karp, and Peter H. Loedel. 2003. "The Euro, Economic Interests and Multi-level Governance: Examining Support for the Common Currency." *European Journal of Political Research* 42:685–703.
- Bryk, Anthony, and Stephen W. Raudenbush. 1992. *Hierarchical Linear Models: Applications and Data Analysis Methods*. Newbury Park, CA: Sage.
- Burden, Barry, and David Kimball. 1998. "A New Approach to the Study of Ticket Splitting." *American Political Science Review* 92:533–544.
- Clarke, Harold D., and Marianne C. Stewart. 1994. "Prospections, Retrospections, and Rationality: The 'Bankers' Model of Presidential Approval Reconsidered." *American Journal of Political Science* 38:1104–1123.
- Cohen, Jeffrey E. 2004. "Economic Perceptions and Executive Approval in Comparative Perspective." *Political Behavior* 26:27–43.
- DeGroot, Morris H., and Mark J. Schervish. 2002. *Probability and Statistics*, 3rd ed. New York: Addison Wesley.
- Efron, Bradley. 1982. *The Jackknife, the Bootstrap and Other Resampling Plans*. Philadelphia, PA: Society for Industrial and Applied Mathematics.
- Greene, William H. 2003. *Econometric Analysis*, 5th ed. Englewood, NJ: Prentice Hall.
- Guerin, Daniel, Jean Crete, and Jean Mercier. 2001. "A Multilevel Analysis of the Determinants of Recycling Behavior in the European Countries." *Social Science Research* 30:195–218.
- Hanushek, Eric A. 1974. "Efficient Estimators for Regressing Regression Coefficients." *American Statistician* 28:66–67.
- Hanushek, Eric A., and John E. Jackson. 1977. *Statistical Methods for Social Scientists*. New York: Academic.
- Herron, Michael C., and Kenneth W. Shotts. 2003. "Using Ecological Inference Point Estimates as Dependent Variables in Second-Stage Linear Regressions." *Political Analysis* 11:44–64.
- Jusko, Karen Long, and W. Phillips Shively. 2005. "Applying a Two-Step Strategy to the Analysis of Cross-National Public Opinion Data." *Political Analysis*. doi:10.1093/pan/mpi030.
- Kaltenthaler, Karl C., and Christopher J. Anderson. 2001. "Europeans and Their Money: Explaining Public Support for the Common European Currency." *European Journal of Political Research* 40:139–170.
- King, Gary. 1997. *A Solution to the Ecological Inference Problem: Reconstructing Individual Behavior from Aggregate Data*. Princeton, NJ: Princeton University Press.
- Long, J. Scott, and Laurie H. Ervin. 2000. "Using Heteroscedasticity Consistent Standard Errors in the Linear Regression Model." *American Statistician* 54:217–224.
- Lubbers, Marcel, Mérove Gijssberts, and Peer Scheepers. 2002. "Extreme Right-Wing Voting in Western Europe." *European Journal of Political Research* 41:345–378.
- MacKinnon, James G., and Halbert White. 1985. "Some Heteroscedasticity Consistent Covariance Matrix Estimators with Improved Finite Sample Properties." *Journal of Econometrics* 29:305–325.
- MacKuen, Michael B., Robert S. Erikson, and James A. Stimson. 1989. "Macropartisanship." *American Political Science Review* 83:1125–1142.
- MacKuen, Michael B., Robert S. Erikson, and James A. Stimson. 1992. "Peasants or Bankers? The American Electorate and the U.S. Economy." *American Political Science Review* 86:597–611.
- Maddala, G. S. 2001. *Introduction to Econometrics*, 3rd ed. West Sussex: John Wiley and Sons.
- Norpoth, Helmut. 1996. "Presidents and the Prospective Voter." *Journal of Politics* 58:776–792.
- Oppenheimer, Bruce. 1996. "The Importance of Elections in a Strong Congressional Party Era." In *Do Elections Matter?* eds. Benjamin Ginsberg and Alan Stone. Armonk, NY: M. E. Sharpe, pp. 120–139.
- Patel, Jagdish K., and Campbell B. Read. 1996. *Handbook of the Normal Distribution*. New York: Marcel Dekker.
- Peffley, Mark, and Robert Rohrschneider. 2003. "Democratization and Political Tolerance in Seventeen Countries: A Multi-level Model of Democratic Learning." *Political Research Quarterly* 56:243–257.
- Rohrschneider, Robert, and Stephen Whitefield. 2004. "Support for Foreign Ownership and Integration in Eastern Europe; Economic Interests, Ideological Commitments, and Democratic Context." *Comparative Political Studies* 37:313–339.
- Saxonhouse, Gary R. 1976. "Estimated Parameters as Dependent Variables." *American Economic Review* 66:178–183.
- Steenbergen, Marco R., and Bradford S. Jones. 2002. "Modeling Multilevel Data Structures." *American Journal of Political Science* 46:218–237.
- Taylor, D. Garth. 1980. "Procedures for Evaluating Trends in Public Opinion." *Public Opinion Quarterly* 44:86–100.
- White, Halbert. 1980. "A Heteroscedastically-Consistent Covariance Matrix Estimator and a Direct Test for the Heteroscedasticity." *Econometrica* 48:817–838.